


Galveston, TX 2014

Jason Schillerstrom, MD schillerstr@uthscsa.edu

Key Points

- Guardianships are either rare or common depending on your perspective.
- A client must be functionally impaired to be incapacitated.
- Executive Function is the strongest cognitive predictor of decisional capacity and self-care abilities.
- Neuropsychological testing *informs* the capacity assessment; it does not replace it.

Scope of the Problem



Scope of the Problem

- How many elders >65yrs in Texas?
- How many 18-64yrs with disability in Texas?
- How many completed in-home APS investigations in FY 2013?
- How many validated in-home APS investigations?
- How many guardianships are there in Texas?

Scope of the problem

- How many elders >65yrs in Texas?
 - 2,954,972
- How many 18-64yrs with disability in Texas?
 - \circ 1,710,430 (elders + disabled = 4,665,402)
- How many completed in-home investigations in FY 2013?
 - 69,383 (=1.5% of elders + disabled)
- How many validated in-home investigations?
 - 48,392
- · How many guardianships are there in Texas?
 - 52,000 (= 1.1% of elders + disabled)

Questions to Answer for APS

- Does the APS client have a psychiatric diagnosis?
- Does the client have decision making capacity?
- What level of care do they need?
- Do they need to be emergently removed from their home?





Find the full text of this and thousands of other resources from leading experts in dozens of legal practice areas in the <u>UT Law CLE eLibrary (utcle.org/elibrary)</u>

Title search: Determination of Capacity

Also available as part of the eCourse

<u>Determination of Capacity; Creating a Guardianship; plus Top 10 Mistakes</u>

First appeared as part of the conference materials for the $16^{\rm th}$ Annual Estate Planning, Guardianship and Elder Law Conference session "Determination of Capacity"